

Western Australia WHEAT STANDARDS 2025/26



These standards are to be applied on individual truck loads only and not averaged over a number of loads in Western Australia.
For South Australian and western Victorian receival standards, visit vitterra.com.au.

CODE		GRADE										
		H1	H2	APW1	APW2	ASW9	AWW1	AUH2	UPW	AWW2	AUW1	FED1
INFRADEC												
PR	Protein (minimum)	13%	11.5%	10.5%	10.0%	9.0%	NO MIN	11.5%	10.5%	NO MIN	10.5%	NO MIN
PR	Protein (maximum)	25.0%	25.0%	25.0%	25.0%	25%	25.0%	25.0%	11.5%	25.0%	25.0%	25.0%
MO	Moisture Content (maximum)	12.5	12.5	12.5	12.5	12.5	13.5	12.5	12.5	13.5	12.5	13.5
TEMP	Temperature (maximum)	50	50	50	50	50	50	50	50	50	50	50
HECTOLITRE WEIGHT (1/2 LITRE)												
TW	Weight (Kg/hl) (minimum)	74	74	74	74	74	74	71	71	68	68	62
TW	Weight (grams) (minimum)	370	370	370	370	370	370	355	355	340	340	310
FOREIGN SEEDS (1/2 LITRE COUNT)												
S1	Type 1 Seeds (maximum) Double gees, Mexican Poppy, Parthenium Weed	8	8	8	8	8	8	8	8	8	8	8
3A	Type 3A Seeds (maximum) Bathurst Burr, Caltrop, Cape Tulip, Thornapple	2	2	2	2	2	2	2	2	2	2	2
3B	Type 3B Seeds (maximum) Vetch (Tare), Vetch (Commercial)	4	4	4	4	4	4	4	4	4	4	4
3C	Type 3C Seeds (maximum) Heliotrope (Blue), Heliotrope (Common)	8	8	8	8	8	8	8	8	8	8	8
S4	Type 4 Seeds (maximum) Bindweed (Field), Paddy Melon, Hexham Scent (Melilotus), Drake Seed Skeleton Weed, Variegated Thistle Nightshades	20	20	20	20	20	20	20	20	20	20	20
S5	Type 5 Seeds (maximum) Patterson's Curse	40	40	40	40	40	40	40	40	40	40	40
S6	Type 6 Seeds (maximum) Saffron Thistle	10	10	10	10	10	10	50	50	50	50	50
7A	Type 7A Seeds (maximum) Chickpeas, Faba Beans, Lupins, Field Peas	1	1	1	1	1	1	10	10	10	10	100
7B	Type 7B Seeds (maximum) Barley, Oats, Wild Oats, Saia Oats, Triticale, Cereal Rye, Bindweed, Turnip Weed, Speargrass and any other seed not mentioned	50	50	50	50	50	50	150	150	150	150	400
EA	Earth (maximum) (Pea size pieces of earth by count)	1	1	1	1	1	1	3	3	3	3	6
SE	Stones Larger than 2mm in diameter or length	0.3g	0.3g	0.3g	0.3g	0.3g	0.3g	0.3g	0.3g	0.3g	0.3g	0.3g
LI	Insects – Large Dead or Alive – Note Pea Weevils (maximum dead only) Rutherglen Bugs, Ladybirds, Grasshoppers, Sitona Weevils, Woodbugs, and any other field insect not damaging to stored grain	3	3	3	3	3	3	3	3	3	3	3
PEAL	Pea Weevils – Alive	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL
SN	Whole Snail Shells (maximum)	1	1	1	1	1	1	1	1	10	10	10
BB	Severely Damaged Grains (maximum) Mould, Bin Burnt & Heat Damaged	1	1	1	1	1	1	1	1	1	1	1
FF	Field Fungi/Black Field Fungi	10	10	10	10	10	10	10	10	10	20	40
SP	Sprouted (maximum)*	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NO LIMIT
FN	Falling Number	300	300	300	300	300	300	250	250	200	250	62
UNMILLABLE MATERIAL (% BY WEIGHT)												
SC	Screenings (maximum %)	5	5	5	5	5	5	10	10	10	25	15
SS	Small Foreign Seeds (maximum %)	0.6	0.6	0.6	0.6	0.6	0.6	1.2	1.2	1.2	1.2	1.2
SA	Unmillable-Whiteheads/Chaff (maximum%)	0.6	0.6	0.6	0.6	0.6	0.6	1.2	1.2	1.2	2.6	2.6
PM	Other Non-Objectionable Material Assessment (% by weight)	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.2
BOTTOM TRAY (1/2 LITRE)												
RE	Ryegrass Ergot (maximum length)	2cm	2cm	2cm	2cm	2cm	2cm	2cm	2cm	2cm	2cm	2cm
CE	Cereal Ergot (maximum) (all cereals except ryegrass ergot)	1	1	1	1	1	1	1	1	1	1	1
LS	Loose Smut (maximum)	3	3	3	3	3	3	3	3	3	3	3
EC	Earcockle (maximum)	10	10	10	10	10	10	10	10	15	15	50
SI	Small Insects Dead or Alive Note Dead Stored Grain Insects Only (maximum)	10	10	10	10	10	10	10	10	10	10	10
SAND ANALYSIS (1/2 LITRE)												
SAND	Sand (maximum)	20	20	20	20	20	50	50	50	50	50	50
WEATHER DAMAGED (1BPM COUNT)												
PF	Pink Grains Only (maximum)	10	10	10	10	10	10	10	10	25	25	25
HS	Head Scab/White Grain Disorder (maximum)	5	5	5	5	5	5	5	5	5	5	5
FS	Stained Grain Only (maximum)	25	25	25	25	25	25	75	75	75	75	250
SP	Sprouted (maximum)*	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NIL	NO LIMIT
GS	Dry Green, Sappy Wheat (maximum)	5	5	5	5	5	5	10	10	25	25	NO LIMIT
XDIST	Distorted Wheat (maximum)	5	5	5	5	5	5	10	10	50	50	NO LIMIT
ID	Insect Damaged (maximum)	5	5	5	5	5	5	5	5	10	10	20